arXiv:1105.4214v1 [math.PR] 21 May 2011

An Inequality Related to Bifractional Brownian Motion

Mikhail Lifshits Ilya Tyurin

St.Petersburg State University, Dept Math.Mech., 198504 Stary Peterhof, Bibliotechnaya pl. 2, Russia
Email: `lifts@mail.rcom.ru`

Moscow State University, Dept Mech.Math., 119991 Moscow, Leninskie gory 1, Russia
Email: `itiurin@gmail.com`

AMS 2000 Subject Classification: 60E15, 60G22

Keywords: bifractional Brownian motion, moment inequalities, Bernstein functions.

Abstract

We prove that for any pair of i.i.d. random variables X, Y with finite moment of order $\alpha \in (0,2]$ it is true that

$$\mathbb{E}|X-Y|^\alpha \le \mathbb{E}|X+Y|^\alpha.$$

Surprisingly, this inequality turns out to be related with bifractional Brownian motion. We extend this result to Bernstein functions and provide some counter-examples.

1 Introduction

Let X, Y be i.i.d. random variables with finite expectations. Then it is true that

$$\mathbb{E}|X-Y| \le \mathbb{E}|X+Y|. \tag{1}$$

Inequality (1) appeared recently in the analytic context (properties of integrable functions) [3]. Since (1) is a nice fact itself and it seems not to be well known in the probabilistic world, it is desirable to search for adequate proofs and consider eventual extensions for it. In particular, is it true that

$$\mathbb{E}|X-Y|^\alpha \le \mathbb{E}|X+Y|^\alpha, \tag{2}$$

provided X and Y are i.i.d. and $\mathbb{E}|X|^\alpha < \infty$?

Proving (1) is a non-trivial exercise for a Probability course. In particular, (1) follows from the identity

$$\mathbb{E}|X+Y| - \mathbb{E}|X-Y| = 2\int_0^\infty \left[\mathbb{P}(X>r) - \mathbb{P}(X<-r)\right]^2 dr.$$

However, we are not aware of a similar elementary approach to (2). In this note, we show how (2) suddenly emerges from some recent advances in the theory of random processes.

2 Main result

We prove the following result.

Theorem 1. *Let $\alpha \in (0,2]$. For any pair of i.i.d. random variables X, Y such that $\mathbb{E}|X|^\alpha < \infty$ it is true that*

$$\mathbb{E}|X-Y|^\alpha \le \mathbb{E}|X+Y|^\alpha.$$

Proof: For the proof we will need the definition of the bifractional Brownian motion introduced by Houdré and Villa in [2]. It is a centered Gaussian process $B^{H,K} = \left(B_t^{H,K}, t \ge 0\right)$ with covariance function

$$R^{H,K}(t,s) = \mathbb{E}\left(B_t^{H,K} B_s^{H,K}\right) = \frac{1}{2^K}\left((t^{2H}+s^{2H})^K - |t-s|^{2HK}\right), \ \ t,s \ge 0.$$

Note that letting $K=1$ yields a usual fractional Brownian motion B^H. Originally, the process was defined for the parameters $H \in (0,1]$ and $K \in (0,1]$. Bardina and Es-Sebaiy [1] proved recently that $B^{H,K}$ exists provided that $(H,K) \in \mathcal{D}$, where

$$\mathcal{D} := \{H, K : 0 < H \le 1, 0 < K \le 2, HK \le 1\}.$$

(Actually a hint for this extension is contained in a preceding work by Lei and Nualart [4] who established an integral representation relating $B^{H,K}$ with fractional Brownian motion B^{HK}).

Now we relate our problem to a particular bifractional Brownian motion. Note that for any $u, v \in \mathbb{R}$ and $\alpha \in (0, 2]$ one has

$$|u+v|^\alpha - |u-v|^\alpha = \begin{cases} (|u|+|v|)^\alpha - ||u|-|v||^\alpha, & uv \geq 0 \\ ||u|-|v||^\alpha - (|u|+|v|)^\alpha, & uv < 0 \end{cases},$$

which can be briefly expressed as

$$((|u|+|v|)^\alpha - ||u|-|v||^\alpha)\, sgn(u)sgn(v) = 2^\alpha R^{1/2,\alpha}(|u|,|v|)sgn(u)sgn(v).$$

Let $\left(B_t^{1/2,\alpha}, t \geq 0\right)$ be a bifractional Brownian motion with $H = 1/2$ and $K = \alpha$. Set

$$Z_\alpha := \int_{-\infty}^{+\infty} B_{|u|}^{1/2,\alpha} sgn(u) P(du),$$

where P stands for the common distribution of X and Y. Then

$$\begin{aligned} 0 \leq Var(Z_\alpha) &= \int_{-\infty}^{+\infty}\int_{-\infty}^{+\infty} R^{1/2,\alpha}(|u|,|v|)sgn(u)sgn(v)P(du)P(dv) \\ &= \int_{-\infty}^{+\infty}\int_{-\infty}^{+\infty} \frac{1}{2^\alpha}(|u+v|^\alpha - |u-v|^\alpha)P(du)P(dv) \\ &= \frac{1}{2^\alpha}[\mathbb{E}|X+Y|^\alpha - \mathbb{E}|X-Y|^\alpha]. \end{aligned}$$

□

3 Further extensions and counterexamples

Inequality (2) trivially extends to the case $\alpha = \infty$ in the following sense. Let

$$\begin{aligned} M &= \sup\{r : \mathbb{P}(X < r) < 1\}; \\ m &= \sup\{r : \mathbb{P}(X < r) = 0\}. \end{aligned}$$

Then

$$||X-Y||_\infty = M - m \leq 2\max\{|M|; |m|\} = ||X+Y||_\infty.$$

However, without further assumptions (2) does not hold in general for $2 < \alpha < \infty$. To see this, let us fix $\alpha \in (2, \infty), c > 0$. For any $M \geq c$ let $q = \frac{c}{M}, p = 1 - q$. Let X_M, Y_M be i.i.d. random variables such that

$$\begin{aligned} \mathbb{P}(X_M = 1) &= \mathbb{P}(Y_M = 1) = p; \\ \mathbb{P}(X_M = -M) &= \mathbb{P}(Y_M = -M) = q. \end{aligned}$$

If $M \geq 1$, then

$$\begin{aligned} & \mathbb{E}|X_M - Y_M|^\alpha - \mathbb{E}|X_M + Y_M|^\alpha \\ = \; & 2pq\left[(M+1)^\alpha - (M-1)^\alpha\right] - 2^\alpha M^\alpha q^2 - 2^\alpha p^2 \\ \geq \; & 4pq\alpha M^{\alpha-1} - 2^\alpha M^\alpha q^2 - 2^\alpha p^2 \\ = \; & M^{\alpha-2}(4p\alpha c - 2^\alpha c^2) - 2^\alpha p^2. \end{aligned}$$

Hence, whenever $c < 2^{2-\alpha}\alpha$ and M is large enough,

$$\mathbb{E}|X_M - Y_M|^\alpha - \mathbb{E}|X_M + Y_M|^\alpha > 0$$

and (2) fails.

It is natural to ask for which class of functions $F(\cdot)$ inequality (2) extends to

$$\mathbb{E}\, F(|X-Y|) \leq \mathbb{E}\, F(|X+Y|). \tag{3}$$

Recall that a function $G : \mathbb{R}_+ \to \mathbb{R}_+$ is called *Bernstein function*, if it admits a representation

$$G(\lambda) = a + b\lambda + \int_0^\infty \left(1 - e^{-t\lambda}\right) \mu(dt),$$

where $a, b \geq 0$, and a measure μ on $(0,\infty)$ satisfies integrability condition $\int_0^\infty \min(t,1)\mu(dt) < \infty$. In a recent monograph [5] one can find an exhaustive information about the importance of Bernstein functions in Analysis and Probability, their properties, as well as a large list of such functions.

Theorem 2. *Let G be a Bernstein function. Then $F(\lambda) := G(\lambda^2)$ satisfies (3) for any pair of i.i.d. random variables X, Y.*

Proof: Since

$$F(\lambda) = a + b\lambda^2 + \int_0^\infty \left(1 - e^{-t\lambda^2}\right) \mu(dt),$$

it is enough to check (3) for elementary functions $F_t(\lambda) := -e^{-t\lambda^2}$. Indeed, from the identity

$$e^{-t|x-y|^2} - e^{-t|x+y|^2} = 2e^{-tx^2-ty^2} \sum_{n=0}^\infty \frac{(2txy)^{2n+1}}{(2n+1)!}$$

we infer that

$$\mathbb{E}\left[F_t(|X+Y|) - F_t(|X-Y|)\right] = 2\sum_{n=0}^\infty \frac{(2t)^{2n+1}}{(2n+1)!} \left[\mathbb{E}\left(X^{2n+1} e^{-tX^2}\right)\right]^2 \geq 0.$$

□

Theorem 1 is a particular case of Theorem 2 because $G(\lambda) = \lambda^{\alpha}$ are Bernstein functions whenever $0 < \alpha < 1$.

Acknowledgement. The authors are grateful to H.Kempka, A. Koldobskii, W. Linde, and A. Nekvinda for setting the problem and for valuable discussions.

The research was supported by the RFBR-DFG grant 09-01-91331, RFBR grants 10-01-00154a, 10-01-00397a, as well as by Federal Focused Programme 2010-1.1-111-128-033.

References

[1] X. Bardina, K. Es-Sebaiy (2011), An extension of bifractional Brownian motion, Commun. Stochast. Analysis, 5, 333–340; `arXiv:1002.3680`.

[2] C. Houdré, J. Villa (2003), An example of infinite dimensional quasi-helix, Contemporary Mathematics, vol. 336, Amer. Math. Soc., pp. 195–201.

[3] H. Kempka, A. Nekvinda (2011), Private communication.

[4] P. Lei, D. Nualart (2009), A decomposition of the bifractional Brownian motion and some applications, Statist. Probab. Letters, 79, 619–624; `arXiv:0803.2227`.

[5] R. Schilling, R. Song, Z. Vondraček (2011), Bernstein Functions. Ser. Studies in Math., vol.37, de Gruyter.